


# 2022 Identification and Treatment of Pediatric Pelvic Fracture Clinical Guideline



# 2022 Identification and Treatment of Pediatric Pelvic Fracture Clinical Guideline

## Vital Signs by Age Group

### Common Causes for Pediatric Pelvis Fracture (Result of High Force)

- MVC
- Pedestrian vs. MVC
- Falls from Significant Height
- Bicycle or Motor Bike Injuries
- High Energy Sports

Age	RR	HR	Systolic BP	Diastolic BP	Weight, kg	HR > 2 SD from Normal Range
Newborn	30-50	120-160	50-70	30-60	2-3	> 180
Infant 1-12 mos	20-30	80-140	70-100	53-66	4-10	> 180
Toddler 1-3 yrs	20-30	80-130	80-110	53-66	10-14	> 140
Preschooler	20-30	80-120	80-110	55-69	14-18	> 140
School Age 6-12 yrs	20-30	70-110	80-120	57-71	20-42	> 130
Adolescent > 13 yrs	12-20	55-105	110-120	66-80	> 50	> 110

# 2022 Identification and Treatment of Pediatric Pelvic Fracture Clinical Guideline

## Unstable Pelvic Fractures and Sheets in a Kocher

Apply pelvic binder or sheet over the greater trochanters of the femur

<https://youtu.be/5bYyySzIMIM>