


2022 Identification and Treatment of Pediatric Pelvic Fracture Clinical Guideline



2022 Identification and Treatment of Pediatric Pelvic Fracture Clinical Guideline

Vital Signs by Age Group

Common Causes for Pediatric Pelvis Fracture (Result of High Force)

- MVC
- Pedestrian vs. MVC
- Falls from Significant Height
- Bicycle or Motor Bike Injuries
- High Energy Sports

Age	RR	HR	Systolic BP	Diastolic BP	Weight, kg	HR > 2 SD from Normal Range
Newborn	30-50	120-160	50-70	30-60	2-3	> 180
Infant 1-12 mos	20-30	80-140	70-100	53-66	4-10	> 180
Toddler 1-3 yrs	20-30	80-130	80-110	53-66	10-14	> 140
Preschooler	20-30	80-120	80-110	55-69	14-18	> 140
School Age 6-12 yrs	20-30	70-110	80-120	57-71	20-42	> 130
Adolescent > 13 yrs	12-20	55-105	110-120	66-80	> 50	> 110

2022 Identification and Treatment of Pediatric Pelvic Fracture Clinical Guideline

Unstable Pelvic Fractures and Sheets in a Kocher

Apply pelvic binder or sheet over the greater trochanters of the femur

<https://youtu.be/5bYyySzIMIM>